

ONDERO

Ondero is a beefsteak tomato with High Resistance to Fusarium wilt and Fusarium crown and root rot. Very good fruit quality and high yield potential.



- High yield potential
- Very good fruit quality throughout the entire crop cycle
- Can be grown un-grafted

Features

- AFW 270 - 300 grams
- High yield potential
- Open, balanced and generative plant type
- Uniform, slightly flat ribbed fruits with attractive color
- Less white parts internally in fruits
- Good fruit firmness and strong against microcracking
- Fast plant development, during the season can produce 1.5 - 2 extra trusses
- Glossy fruits

Advantages

- High yield potential
- Glossy fruits
- Good overall fruit quality

Agronomic Tips

- Grafting on DR0141TX, Maxifort in long crops in warm regions is recommended. Fruit fall: avoid high humidity associated to heat stress during the low light season by properly managing the heating system and maintaining active climate. Under the same conditions, irrigation and EC may interfere negatively in the incidence of fruit fall. Regular leaf removal on time does help minimize fruit fall and quality issues. Planting after 10 January to avoid low light levels during setting and development of first trusses.

Disease Resistance

HR **ToMV:0-2/ToTV/Pf:A-E/Fol:1,2/For**

IR **On**

Disease Resistance Key

- **Fol = Fusarium wilt**
- **For = Fusarium crown and root rot**
- **On = Powdery mildew**
- **ToMV = Tomato mosaic**
- **ToTV = Tomato torrado**

Characteristics

| | |
|----------------------|-------------------------|
| BRIX POTENTIAL | 4 - 5 |
| SEGMENT | XL |
| SIZE | Beef |
| EXTERNAL COLOR | Red |
| SHAPE | Round |
| AVERAGE WEIGHT | 270-300 grams |
| HARVEST METHOD | Loose |
| HARVEST UNIFORMITY | Good Uniformity |
| HARVESTING SEASON | All YearLit Crop |
| JOINTLESS | No |
| PLANT TYPE | Indeterminate |
| VIGOR | Strong |
| RELATIVE MATURITY | Medium |
| SOWING SEASON | All YearLit Crop |
| TRANSPLANTING SEASON | All YearLit Crop |

HR = High Resistance. IR = Intermediate Resistance. To find out more about disease resistance and the applicable levels of disease resistance, https://www.worldseed.org/wp-content/uploads/2017/05/Definition_on_reaction_plants_to_pests_2017_final.pdf. To see the Disease Resistance Abbreviation Guide please visit <https://www.vegetables.bayer.com/ca/en-ca/resources/disease-code-abbreviations.html>

Performance may vary from location to location and from year to year, as local growing, soil and climatic conditions may vary. Growers should evaluate data from multiple locations and years whenever possible and should consider the impacts of these conditions on the grower's operation Any recommendations provided on this material are based upon observations and feedback received from a limited number of trials and geographies. These recommendations should be considered as one reference point and should not be substituted for the professional opinion of agronomists, entomologists or other relevant experts evaluating specific conditions. De Ruiter® is a registered trademark of Bayer Group. All other trademarks displayed on this material are either registered or unregistered trademarks of Bayer Group, one of its related affiliates or others and used by permission. © 2004 – 2023 Bayer Group. All rights reserved.